



Nobbs, A. H. (2016). Getting to the heart of the matter: Role of *Streptococcus mutans* adhesin Cnm in systemic disease. *Virulence*, 8(1), 1-4. <https://doi.org/10.1080/21505594.2016.1212157>

Peer reviewed version

Link to published version (if available):  
[10.1080/21505594.2016.1212157](https://doi.org/10.1080/21505594.2016.1212157)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Taylor & Francis at <http://www.tandfonline.com/doi/abs/10.1080/21505594.2016.1212157>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

1     **Getting to the heart of the matter: role of *Streptococcus mutans* adhesin Cnm in systemic disease**

2     Angela Nobbs

3     School of Oral and Dental Sciences, University of Bristol, Bristol, UK

4

5     **Keywords:** *Streptococcus*; infective endocarditis; adhesin; collagen-binding protein; *Lactococcus*

6     *lactis*

7

8     *Streptococcus mutans* is one of an estimated 700 prokaryote species that are recognised as  
9     constituents of the oral microbiota <sup>1</sup>. *S. mutans* is exclusively found as a member of the  
10    polymicrobial biofilm communities that comprise dental plaque <sup>2</sup>, and is perhaps most notorious as  
11    the first bacterium to be identified as a major aetiological agent of dental caries (tooth decay) <sup>3</sup>. The  
12    incidence of this disease has declined with the introduction of population-based prevention  
13    measures <sup>4</sup>. Nonetheless, dental caries remains one of the most ubiquitous bacterial infections of  
14    humans and represents a significant financial burden to the healthcare system <sup>5, 6</sup>.

15

16    *S. mutans* and several other streptococcal species that commonly inhabit the oral cavity are  
17    collectively known as the ‘viridans group’ streptococci and together can comprise up to 80% of early  
18    dental plaque <sup>7, 8</sup>. Outside of the oral niche, however, this group of bacteria is also particularly  
19    recognised for its association with heart condition infective endocarditis (IE). Together with  
20    staphylococci and enterococci, the viridans group streptococci account for 80-90% of all IE cases <sup>9</sup>.  
21    Whilst relatively rare (3-10 cases per 100,000 individuals), the one year mortality rate for this  
22    infection of the heart valves remains at ca. 30%, with treatment options frequently incorporating  
23    surgery and long-term administration of antibiotics <sup>10</sup>. Given the current crisis of increasing incidence  
24    in antibiotic resistance within the global microbial population <sup>11</sup>, there is considerable pressure to  
25    devise alternative treatment strategies for IE. To achieve this, however, greater understanding of the  
26    pathogenic mechanisms that underpin this disease is required.

27

28 For the viridans group streptococci, the initial step in IE pathogenesis is bacterial entry into the  
29 bloodstream. Such transient bacteraemias arise following disruption of the oral mucosae, often  
30 simply as a result of daily practices (e.g. toothbrushing, flossing)<sup>12-15</sup>. Upon transiting from the oral  
31 cavity to the cardiovascular system, these streptococci must then adhere to the endothelium of the  
32 heart valves, where they promote deposition of fibrin and blood platelets to form an infective  
33 vegetation (clot)<sup>9, 16, 17</sup>. In this issue, Freires et al.<sup>18</sup> utilise a surrogate host expression system to  
34 demonstrate that *S. mutans* surface adhesin Cnm (collagen-binding protein of *S. mutans*) is essential  
35 for this process.

36

37 Strains of *S. mutans* can be divided across four capsular polysaccharide serotypes (*c*, *e*, *f* and *k*), of  
38 which serotype *c* is the most prevalent within the oral niche<sup>19, 20</sup>. Serotypes *e*, *f* and *k* have been  
39 found to express Cnm or the closely-related collagen-binding protein Cbm. By contrast, serotype *c*  
40 strains, which comprise ca. 75% of isolates, typically lack the *cnm* locus<sup>21-24</sup>. Intriguingly, the less  
41 abundant serotypes are highly overrepresented among isolates associated with *S. mutans* extra-oral  
42 infections<sup>25, 26</sup>. This disparity in distribution provided some of the first evidence that such collagen-  
43 binding adhesins may make a critical contribution to the capacity for *S. mutans* to cause systemic  
44 disease. Cnm has been shown to promote attachment to extracellular matrix (ECM) proteins  
45 collagen (types I, II, III and IV) and laminin<sup>21, 25, 27-29</sup>, and to facilitate invasion of cardiac endothelial  
46 cells<sup>28, 30</sup>. The role of Cnm as a potential virulence factor was also demonstrated using the *Galleria*  
47 *mellonella* wax worm model of systemic infection<sup>28, 30</sup>. Such studies were primarily performed using  
48 *S. mutans*  $\Delta cnm$  knockout mutants and corresponding complemented strains. However,  
49 *Streptococcus* bacteria are notorious for exhibiting adhesin redundancy<sup>31-33</sup>. This feature likely  
50 contributes to the overwhelming success of these bacteria as host colonisers, but can make it  
51 challenging to unequivocally ascribe an adhesive function(s) to a specific protein. One way to  
52 address this issue is to utilise a heterologous expression system and a successful strain for which

53 there is precedent with *Streptococcus* proteins is *Lactococcus lactis*<sup>34-38</sup>. As a Gram-positive coccus,  
54 *L. lactis* shares many of the systems required for surface protein export and display and yet, as a  
55 dairy industry starter microorganism, it lacks capacity to interact strongly with human cells and  
56 tissues<sup>39</sup>. Consequently, *L. lactis* can serve as an excellent 'blank canvas' with which gain of function  
57 can be explored following expression of a heterologous protein.

58

59 Using *L. lactis* expressing Cnm, Freires et al.<sup>18</sup> demonstrate unambiguously that this adhesin  
60 mediates adhesion to ECM components collagen type I and laminin, by both direct whole cell  
61 binding assays and complementary inhibition studies using anti-Cnm serum. Cnm is also shown to  
62 confer capacity to invade human coronary artery endothelial cells (HCAEC), and Cnm<sup>+</sup> *L. lactis*  
63 exhibits significantly enhanced virulence using the *G. mellonella* model of systemic disease  
64 compared to parent strain. Additionally, in contrast to parent strain, *L. lactis* expressing Cnm is able  
65 to bind freshly extirpated human aortic valve tissue. Evidence suggests that one way IE might be  
66 initiated is through adherence of viridans group streptococci to exposed ECM proteins of damaged  
67 heart valves<sup>40</sup>. The SEM images shown in Freires et al.<sup>18</sup> support this mechanism, with bacterial  
68 adhesion to collagenous fibrils present in areas of damage clearly visible. Nonetheless, this study  
69 also presents examples of binding to supposedly intact endothelium. Such observations are of  
70 interest, as they imply that Cnm may also facilitate recognition of endothelial receptors. Direct  
71 binding to endothelial cell lines in vitro has been demonstrated previously for *Streptococcus* bacteria  
72<sup>41-44</sup>, but this has yet to be considered as a potential mechanism in IE. Such a possibility is worthy of  
73 investigation in future studies.

74

75 Evidence of a role for specific streptococcal adhesins in IE has proven difficult to obtain using animal  
76 models, possibly reflecting the challenge posed by adhesin redundancy and/or the capacity for  
77 bacteria to utilise multiple mechanisms. The most striking example of this perhaps was seen for  
78 viridans group member *Streptococcus sanguinis*, for which no single deletion of any of its 33 surface

79 (LPxTG-anchored) proteins significantly affected IE outcome<sup>45</sup>. Again, this is where a surrogate host  
80 can offer advantages. Using a rabbit model of IE in which the animals are co-inoculated with both  
81 parent and Cnm<sup>+</sup> *L. lactis*, Freires et al.<sup>18</sup> show that Cnm confers a 67% increase in infectivity. This  
82 serves to reinforce the proposed role of Cnm as enabling initial contact and retention of *S. mutans*  
83 with valve endocardium.

84

85 While surface expression of heterologous proteins can be successfully achieved with *L. lactis*, one  
86 aspect that may not be faithfully reproduced is with posttranslational modifications such as  
87 glycosylation. This can, however, in itself be informative. In *S. mutans* Cnm has been shown to be co-  
88 transcribed with GT-A type glycosyltransferase PgfS, which appears to modify Cnm through *O*-  
89 glycosylation of its threonine-rich B domain<sup>46</sup>. Freires et al.<sup>18</sup> show that this glycosylation does not  
90 occur in *L. lactis*, resulting in expression of a lower MW variant of Cnm with greater susceptibility to  
91 proteinase K degradation. Since Cnm<sup>+</sup> *L. lactis* exhibits significant interactions with ECM proteins and  
92 cardiac endothelium, these data indicate that it is the protein backbone rather than the sugar  
93 modifications of Cnm that mediates its adhesive properties. Nonetheless, this study also provides  
94 evidence that the stability conferred by *O*-glycosylation may be critical for Cnm-mediated adhesion  
95 in vivo. Deciphering the precise contribution that *O*-glycosylation makes to the overall functionality  
96 of Cnm in adhesion and pathogenesis again represent important areas for future research.

97

98 It is becoming increasingly evident that a diverse array of mechanisms are utilised by different  
99 members of the viridans group streptococci to promote thrombosis and the progression of IE.

100 Studies such as these of Freires et al.<sup>18</sup> are helping to advance understanding of this complex host-  
101 microbe interplay and the development of new anti-infection strategies that might help move away  
102 from the current complete reliance on antibiotics.

103

104 **Disclosure of potential conflicts of interest**

No potential conflicts of interest were disclosed.

## References

1. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, Lakshmanan A, Wade WG. The human oral microbiome. *J Bacteriol* 2010; 192:5002-17.
2. Marsh PD. Microbial ecology of dental plaque and its significance in health and disease. *Adv Dent Res* 1994; 8:263-71.
3. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* 1986; 50:353-80.
4. Iheozor-Ejiofor Z, Worthington HV, Walsh T, O'Malley L, Clarkson JE, Macey R, Alam R, Tugwell P, Welch V, Glenny AM. Water fluoridation for the prevention of dental caries. *Cochrane Database Syst Rev* 2015:Cd010856.
5. Bagramian RA, Garcia-Godoy F, Volpe AR. The global increase in dental caries. A pending public health crisis. *Am J Dent* 2009; 22:3-8.
6. Collaborators GBoDS. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: A systematic analysis for the global burden of disease study 2013. *Lancet* 2015; 386:743-800.
7. Diaz PI, Chalmers NI, Rickard AH, Kong C, Milburn CL, Palmer RJ, Jr., Kolenbrander PE. Molecular characterization of subject-specific oral microflora during initial colonization of enamel. *Appl Environ Microbiol* 2006; 72:2837-48.
8. Dige I, Nyengaard JR, Kilian M, Nyvad B. Application of stereological principles for quantification of bacteria in intact dental biofilms. *Oral Microbiol Immunol* 2009; 24:69-75.
9. Que YA, Moreillon P. Infective endocarditis. *Nat Rev Cardiol* 2011; 8:322-36.
10. Cahill TJ, Prendergast BD. Current controversies in infective endocarditis. *F1000Res* 2015; 4.
11. Premanandh J, Samara BS, Mazen AN. Race against antimicrobial resistance requires coordinated action - an overview. *Front Microbiol* 2015; 6:1536.

- 131 12. Crasta K, Daly CG, Mitchell D, Curtis B, Stewart D, Heitz-Mayfield LJ. Bacteraemia due to dental  
132 flossing. *J Clin Periodontol* 2009; 36:323-32.
- 133 13. Zhang W, Daly CG, Mitchell D, Curtis B. Incidence and magnitude of bacteraemia caused by  
134 flossing and by scaling and root planing. *J Clin Periodontol* 2013; 40:41-52.
- 135 14. Mougeot FK, Saunders SE, Brennan MT, Lockhart PB. Associations between bacteremia from  
136 oral sources and distant-site infections: Tooth brushing versus single tooth extraction. *Oral Surg*  
137 *Oral Med Oral Pathol Oral Radiol* 2015; 119:430-5.
- 138 15. Lockhart PB, Brennan MT, Sasser HC, Fox PC, Paster BJ, Bahrani-Mougeot FK. Bacteremia  
139 associated with toothbrushing and dental extraction. *Circulation* 2008; 117:3118-25.
- 140 16. Keane C, Petersen HJ, Tilley DO, Haworth J, Cox D, Jenkinson HF, Kerrigan SW. Multiple sites on  
141 *Streptococcus gordonii* surface protein PadA bind to platelet GPIIb/IIIa. *Thromb Haemost* 2013;  
142 110:1278-87.
- 143 17. Kerrigan SW. The expanding field of platelet-bacterial interconnections. *Platelets* 2015; 26:293-  
144 301.
- 145 18. Freires IA, Aviles-Reyes A, Kitten T, Simpson-Haidaris PJ, Swartz M, Knight PA, Rosalen PL, Lemos  
146 JA, Abranches J. Heterologous expression of *Streptococcus mutans* Cnm in *Lactococcus lactis*  
147 promotes intracellular invasion, adhesion to human cardiac tissues and virulence. *Virulence*  
148 2016:1-12.
- 149 19. Nakano K, Ooshima T. Serotype classification of *Streptococcus mutans* and its detection outside  
150 the oral cavity. *Future Microbiol* 2009; 4:891-902.
- 151 20. Shibata Y, Ozaki K, Seki M, Kawato T, Tanaka H, Nakano Y, Yamashita Y. Analysis of loci required  
152 for determination of serotype antigenicity in *Streptococcus mutans* and its clinical utilization. *J*  
153 *Clin Microbiol* 2003; 41:4107-12.
- 154 21. Lapirattanakul J, Nakano K, Nomura R, Leelataweewud P, Chalermarp N, Klaophimai A,  
155 Srisatjaluk R, Hamada S, Ooshima T. Multilocus sequence typing analysis of *Streptococcus*

156 *mutans* strains with the *cnm* gene encoding collagen-binding adhesin. J Med Microbiol 2011;  
 157 60:1677-84.

158 22. Lapirottanakul J, Nomura R, Nemoto H, Naka S, Ooshima T, Nakano K. Multilocus sequence  
 159 typing of *Streptococcus mutans* strains with the *cbm* gene encoding a novel collagen-binding  
 160 protein. Arch Oral Biol 2013; 58:989-96.

161 23. Nakano K, Lapirottanakul J, Nomura R, Nemoto H, Alaluusua S, Gronroos L, Vaara M, Hamada S,  
 162 Ooshima T, Nakagawa I. *Streptococcus mutans* clonal variation revealed by multilocus sequence  
 163 typing. J Clin Microbiol 2007; 45:2616-25.

164 24. Nomura R, Nakano K, Naka S, Nemoto H, Masuda K, Lapirottanakul J, Alaluusua S, Matsumoto  
 165 M, Kawabata S, Ooshima T. Identification and characterization of a collagen-binding protein,  
 166 Cbm, in *Streptococcus mutans*. Mol Oral Microbiol 2012; 27:308-23.

167 25. Nakano K, Nomura R, Taniguchi N, Lapirottanakul J, Kojima A, Naka S, Senawongse P, Srisatjaluk  
 168 R, Gronroos L, Alaluusua S, et al. Molecular characterization of *Streptococcus mutans* strains  
 169 containing the *cnm* gene encoding a collagen-binding adhesin. Arch Oral Biol 2010; 55:34-9.

170 26. Nakano K, Nemoto H, Nomura R, Homma H, Yoshioka H, Shudo Y, Hata H, Toda K, Taniguchi K,  
 171 Amano A, et al. Serotype distribution of *Streptococcus mutans* a pathogen of dental caries in  
 172 cardiovascular specimens from Japanese patients. J Med Microbiol 2007; 56:551-6.

173 27. Sato Y, Okamoto K, Kagami A, Yamamoto Y, Igarashi T, Kizaki H. *Streptococcus mutans* strains  
 174 harboring collagen-binding adhesin. J Dent Res 2004; 83:534-9.

175 28. Abranches J, Miller JH, Martinez AR, Simpson-Haidaris PJ, Burne RA, Lemos JA. The collagen-  
 176 binding protein Cnm is required for *Streptococcus mutans* adherence to and intracellular  
 177 invasion of human coronary artery endothelial cells. Infect Immun 2011; 79:2277-84.

178 29. Nomura R, Naka S, Nemoto H, Inagaki S, Taniguchi K, Ooshima T, Nakano K. Potential  
 179 involvement of collagen-binding proteins of *Streptococcus mutans* in infective endocarditis. Oral  
 180 Dis 2013; 19:387-93.



- 181 30. Aviles-Reyes A, Miller JH, Simpson-Haidaris PJ, Lemos JA, Abranches J. Cnm is a major virulence  
182 factor of invasive *Streptococcus mutans* and part of a conserved three-gene locus. Mol Oral  
183 Microbiol 2014; 29:11-23.
- 184 31. Nobbs AH, Lamont RJ, Jenkinson HF. *Streptococcus* adherence and colonization. Microbiol Mol  
185 Biol Rev 2009; 73:407-50.
- 186 32. Zhang Y, Lei Y, Nobbs A, Khammanivong A, Herzberg MC. Inactivation of *Streptococcus gordonii*  
187 SspAB alters expression of multiple adhesin genes. Infect Immun 2005; 73:3351-7.
- 188 33. Jakubovics NS, Brittan JL, Dutton LC, Jenkinson HF. Multiple adhesin proteins on the cell surface  
189 of *Streptococcus gordonii* are involved in adhesion to human fibronectin. Microbiology 2009;  
190 155:3572-80.
- 191 34. Brittan JL, Nobbs AH. Group B *Streptococcus* pili mediate adherence to salivary glycoproteins.  
192 Microbes Infect 2015; 17:360-8.
- 193 35. Asmat TM, Klingbeil K, Jensch I, Burchhardt G, Hammerschmidt S. Heterologous expression of  
194 pneumococcal virulence factor PspC on the surface of *Lactococcus lactis* confers adhesive  
195 properties. Microbiology 2012; 158:771-80.
- 196 36. Holmes AR, Gilbert C, Wells JM, Jenkinson HF. Binding properties of *Streptococcus gordonii* SspA  
197 and SspB (antigen I/II family) polypeptides expressed on the cell surface of *Lactococcus lactis*  
198 MG1363. Infect Immun 1998; 66:4633-9.
- 199 37. Morita C, Sumioka R, Nakata M, Okahashi N, Wada S, Yamashiro T, Hayashi M, Hamada S,  
200 Sumitomo T, Kawabata S. Cell wall-anchored nuclease of *Streptococcus sanguinis* contributes to  
201 escape from neutrophil extracellular trap-mediated bacteriocidal activity. PLoS One 2014;  
202 9:e103125.
- 203 38. Maddocks SE, Wright CJ, Nobbs AH, Brittan JL, Franklin L, Stromberg N, Kadioglu A, Jepson MA,  
204 Jenkinson HF. *Streptococcus pyogenes* antigen I/II-family polypeptide AspA shows differential  
205 ligand-binding properties and mediates biofilm formation. Mol Microbiol 2011; 81:1034-49.

- 206 39. Villatoro-Hernandez J, Kuipers OP, Saucedo-Cardenas O, Montes-de-Oca-Luna R. Heterologous  
207 protein expression by *Lactococcus lactis*. Methods Mol Biol 2012; 824:155-65.
- 208 40. Moreillon P, Que YA. Infective endocarditis. Lancet 2004; 363:139-49.
- 209 41. Stinson MW, Alder S, Kumar S. Invasion and killing of human endothelial cells by viridans group  
210 streptococci. Infect Immun 2003; 71:2365-72.
- 211 42. Vacca-Smith AM, Jones CA, Levine MJ, Stinson MW. Glucosyltransferase mediates adhesion of  
212 *Streptococcus gordonii* to human endothelial cells in vitro. Infect Immun 1994; 62:2187-94.
- 213 43. Vernier A, Diab M, Soell M, Haan-Archipoff G, Beretz A, Wachsmann D, Klein JP. Cytokine  
214 production by human epithelial and endothelial cells following exposure to oral viridans  
215 streptococci involves lectin interactions between bacteria and cell surface receptors. Infect  
216 Immun 1996; 64:3016-22.
- 217 44. Vollmer T, Hinse D, Kleesiek K, Dreier J. Interactions between endocarditis-derived  
218 *Streptococcus gallolyticus* subsp. *gallolyticus* isolates and human endothelial cells. BMC  
219 Microbiol 2010; 10:78.
- 220 45. Turner LS, Kanamoto T, Unoki T, Munro CL, Wu H, Kitten T. Comprehensive evaluation of  
221 *Streptococcus sanguinis* cell wall-anchored proteins in early infective endocarditis. Infect Immun  
222 2009; 77:4966-75.
- 223 46. Aviles-Reyes A, Miller JH, Simpson-Haidaris PJ, Hagen FK, Abranches J, Lemos JA. Modification of  
224 *Streptococcus mutans* Cnm by PgfS contributes to adhesion, endothelial cell invasion, and  
225 virulence. J Bacteriol 2014; 196:2789-97.